




ANTIBACTERIAL DRESSING

MEDIHONEY®

ANTIBACTERIAL HONEY 3-PLY TULLE DRESSING



FEATURES

- Antibacterial *Leptospermum* (Manuka) Honey™ impregnated into a strong 3-ply woven dressing
- Sterile, non-adherent wound dressing

INDICATIONS FOR USE

- Leg/foot ulcers
- Pressure ulcers
- Infected wounds
- Sloughy wounds
- Necrotic wounds
- Malodorous wounds
- Donor and recipient graft sites
- Burns
- Surgical wounds
- Diabetic wounds
- Superficial wounds such as cuts, scratches and abrasions

SUPERIOR WOUND BED PROTECTION

MEDIHONEY® Antibacterial Honey 3-Ply Tulle Dressing protects the wound by creating a barrier against wound pathogens including antibiotic resistant strains therefore reducing the risk of infection. The osmotic action produces an outflow of body fluid which assists the removal of wound bacteria, endotoxins, debris and slough; providing a cleaner wound, rapidly removing malodour and helping to reduce the inflammatory response, oedema and exudate levels. Granulation and epithelialisation are enhanced through provision of the optimal healing environment.

MEDIHONEY® Antibacterial Honey 3-Ply Tulle Dressing creates a micro-environment that supports healing and provides an antibacterial barrier.

international.dermasciences.com

MEDIHONEY® is a trademark of Derma Sciences, Inc.
2017 Derma Sciences, Inc. All rights reserved. MH-TULLE_002E




Derma Sciences, Inc.
104 Shorting Road
Toronto, ON M1S 3S4
P: 416.299.4003 Toll Free: 800.387.5302
Email: customerservice@dermasciences.com



ANTIBACTERIAL DRESSING
MEDIHONEY®

ANTIBACTERIAL HONEY 3-PLY TULLE DRESSING



ITEM#	SIZE	PKG/UNIT
DUP797	2inx2in (5cmx5cm)	5/box
DUP796	4inx4in (10cmx10cm)	5/box

OBSERVATIONS

The wound may appear larger in size and depth as dead tissue is cleaned away. This increase should be followed by an improved appearance of the wound as it heals.

DIRECTIONS FOR USE

1. If required, protect the wound edges with MEDIHONEY® Barrier Cream.
2. Place the dressing onto the wound surface ensuring that the dressing is in contact with the wound bed. Dressings can be placed alongside or on top of one another to cover large or deep areas. Alternatively, the wound may be filled with MEDIHONEY® Antibacterial Medical Honey® before applying the dressing. The dressing may be cut with sterile scissors or folded to the shape of the wound.
3. A suitable secondary sterile dressing should be applied to cover the MEDIHONEY® Antibacterial Honey 3-Ply Tulle Dressing. The secondary dressing should be sufficiently absorbent to manage wound exudate.
4. The wound may be cleaned between dressing changes in accordance with standard wound cleaning protocol.

Suitable for use under compression dressings.

FREQUENCY OF CHANGE

- MEDIHONEY® Antibacterial Honey 3-Ply Tulle Dressing requires changing when they have been diluted by wound exudate and/or when the dressing has gelled. The dressing may be left in place for up to 7 days depending on the amount of wound exudate.
- MEDIHONEY® Antibacterial Honey 3-Ply Tulle Dressing is non-adherent. It is easy to remove without disrupting the wound bed and new tissue.
- Heavily exudating wounds require close monitoring and may require daily dressing changes.

CONTRAINDICATIONS

Do not use on individuals who have a known sensitivity to any component present in the dressing.

PRECAUTIONS

- Transient stinging may be experienced on application of the dressing. This is due to the natural properties of the honey and is not detrimental to the wound. If transient stinging persists and cannot be managed with an analgesic after 30 minutes, remove the dressing and wash the affected area.
- When used on dry wounds, it may be helpful to moisten the dressing to aid removal.
- Use with caution on heavily bleeding wounds.

Pair MEDIHONEY® with our super absorbent cover dressing, XTRASORB®. It's osmotic gradient pulls exudate to the back of the dressing and converts it into a gel, locking it away - even under compression!

